arXiv:1003.0954v1 [cond-mat.str-el] 4 Mar 2010

Quantum critical surface of the zigzag spin chain under magnetic field: Application to superconducting quantum dots

Meihua Chen[1], Sujit Sarkar[2], C. D. Hu[1]
[1]*Physics Department, National Taiwan University, Taipei, R. O. C and*
[2]*Poornaprajna Institute of Scientific Research, 4 Sadashivanagar, Bangalore- 5600 80, India*

(Dated: March 4, 2010)

We analyze the exact ground state of XXZ zigzag spin chain with applied magnetic field and find the quantum critical surface. Using the theorem of positive semi-definite matrix, we can prove that the ground states for a specific region, are fully polarized state and one magnon states. With Bethe ansatz, we argue that this is the quantum critical surface in all cases. A first in the literature, we derive the analytical expression of quantum critical surface for superconducting quantum dots array in the presence of gate voltage.

PACS numbers:

Low dimensional spin systems has been an active field of research [1] for many year. Bethe ansatz has been used widely to solve many new families of integrable Hamiltonians of one dimensional systems [2–5]. A quantum inverse scattering method based on Bethe ansatz has also been developed for dealing with the two-dimensional models [6]. Another method which also enables one to find the exact solutions of various models is the matrix product method [7–10] . For example, it is related to the Affleck-Kennedy-Lieb-Tasaki(AKLT) state [9, 11]. In this work we are going to apply these two methods to the zigzag spin chain model.

A zigzag spin chain is an one dimensional system of spins with exchange interactions with their nearest (NN) and next-nearest neighbors (NNN), denoted by J_1 and J_2 respectively. It is a frustrated system if $J_1 J_2 < 0$ or both of them are positive. In a zigzag spin chain the situation is not clear. Majumdar and Ghosh [12] found that the degenerate dimer states in which neighboring sites form singlets (MG state) are the exact ground states at $\frac{J_1}{J_2} = 2$. At $\frac{J_1}{J_2} = -4$, Hanada et.al. [13] found that the uniformly distributed resonate valence bond (UDRVB) state and the fully polarized (FP) state are degenerate ground states. There are also a few other important theoretical works in zigzag spin chain [14–17]. The zigzag spin chain can be realized in physical systems, such as the oxides, Li_2CuO_2 (multiferroic) [18] and $SrCuO_2$ [19]. In this letter, we find a quantum critical surface on which the FP state is degenerate with the one-magnon states, along with an application to superconducting quantum dots system. Our calculation is based on the concept that the local ground states in a matrix product form can be extended to global ground states.

Model Hamiltonian: We begin with the Hamiltonian of a zigzag spin chain with applied magnetic field,

$$H = \sum_i J_1[\Delta_1 s^z_i s^z_{i+1} + \frac{1}{2}(s^+_i s^-_{i+1} + s^-_i s^+_{i+1})]$$
$$+ \sum_i J_2[\Delta_2 s^z_i s^z_{i+2} + \frac{1}{2}(s^+_i s^-_{i+2} + s^-_i s^+_{i+2})] + B\sum_i s^z_i$$

where J_1 and J_2 are the NN and, NNN interaction, $\Delta_1(\Delta_2)$ is the anisotropy in z axes for NN(NNN) interaction and i is the label of the lattice site. Furthermore, we use the periodic boundary condition and hence we have the coupling between $\vec{s}_{N-1}, \vec{s}_N$ and $\vec{s}_1, \vec{s}_2$. We can dissect the Hamiltonian into many local Hamiltonians. Each local Hamiltonian contains three spins. Thus, the original Hamiltonian can be written as

$$H = \sum_i h_{i,i+1,i+2} = \sum_i (h^0_{i,i+1,i+2} + h^{mag}_{i,i+1,i+2}). \quad (1)$$

where $h^0_{i,i+1,i+2} = \frac{f}{2}[\Delta_1(s^z_i + s^z_{i+2})s^z_{i+1} + \frac{1}{2}[s^-_{i+1}(s^+_i + s^+_{i+2}) + h.c.] + [\Delta_2 s^z_i s^z_{i+2} + \frac{1}{2}(s^+_i s^-_{i+2} + s^-_i s^+_{i+2})]$, and $h^{mag}_{i,i+1,i+2} = \frac{b}{2+x}(s^z_i + x s^z_{i+1} + s^z_{i+2})$. $h^0_{i,i+1,i+2}$ is the local Hamiltonian of exchange interaction, and $h^{mag}_{i,i+1,i+2}$ is the that of applied magnetic field. We have set $b = \frac{B}{J_2}, f = \frac{J_1}{J_2}$ and made scaling $J_2 = 1$. In h^{mag} we have introduced a free parameter x to facilitate later calculation. Note that the Hamiltonian is independent of the value of x if we apply the periodic boundary condition.

The quantum critical surface from local site Hamiltonian: The local Hamiltonian h_{123}, which serves as our starting point, can be diagonalize and the eigenvalues in terms of parameters x, $\Delta_{1,2}$ and f are

$$E_0 = \frac{1}{4}(-2b + \Delta_2 + \Delta_1 f),$$
$$E_1 = \frac{1}{4}(2b + \Delta_2 + \Delta_1 f),$$
$$E_2 = \frac{1}{4}(-2 - 2b - \Delta_2) + \frac{b}{2+x},$$
$$E_3 = \frac{1}{4}(-2 - \Delta_2 + \frac{2bx}{2+x}),$$
$$E_{4\pm} = \frac{4b - (-2 + \Delta_1 f)(x+2) \pm \Gamma_+}{8(x+2)},$$
$$E_{5\pm} = \frac{-4b - (-2 + \Delta_1 f)(x+2) \pm \Gamma_-}{8(x+2)},$$

where

$$\Gamma_{\pm}^2 = [16b^2(-1+x)^2 \pm 8b(2-2\Delta_2+\Delta_1 f)(-2+x+x^2) + 4(-1+\Delta_2)^2 - 4\Delta_1(-1+\Delta_2)f + (8+\Delta_1^2)f^2]$$

We found that $E_0(E_1)$ corresponds to the state $|\downarrow\downarrow\downarrow\rangle\,(|\uparrow\uparrow\uparrow\rangle)$, E_2 and $E_{5\pm}$ correspond to one spin-up states and E_3 and $E_{4\pm}$ to the one spin-down states.

In order to see quantum critical points for entire system, we seek in the spectrum the level crossing between FP state $|\downarrow\downarrow\downarrow\rangle$ and other eigen state. We need at least two other states because for an infinite spin chain with space inversion symmetry any other states must be two-fold degenerate. Inspecting the eigenvalues, we found that E_0, E_2 and E_{5-} are three lowest possible energies. By requiring $E_0 = E_2 = E_{5-}$, we get the following relations $x = \frac{f(f+4\Delta_1)}{2(2+2\Delta_2+\Delta_1 f)}$, $b = \frac{1}{4}(2+2\Delta_2+\Delta_1 f))(2+x)$ or for indication of the quantum critical surface

$$b = 1 + \Delta_2 + \Delta_1 f + \frac{f^2}{8}. \tag{2}$$

The correspond energy of these three degenerate state is

$$E_0 = \frac{-2-\Delta_2-\Delta_1 f}{4} - \frac{f^2}{16} \tag{3}$$

Matrix formulation of the global eigen state: Here, we generalize the results of local Hamiltonian for arbitrary number of spins. The global Hamiltonian of N spins can be obtained by the direct sum of local Hamiltonian.

$$H = h \otimes \hat{1}_{2^{N-3}} \oplus \hat{1}_{2^1} \otimes h \otimes \hat{1}_{2^{N-4}} \oplus \hat{1}_{2^2} \otimes h \otimes \hat{1}_{2^{N-5}} \oplus \cdots \oplus \hat{1}_{2^{N-4}} \otimes h \otimes \hat{1}_{2^1} \oplus \hat{1}_{2^{N-3}} \otimes h,$$

where each basic element h is just the local Hamiltonian $h_{i,i+1,i+2}$ and $\hat{1}_M$ is the identity matrix of rank M. The eigen states of an infinite spin chain can be also found by adding more spins to eigen states of the local Hamiltonian. Consider the matrix

$$m_j = \begin{pmatrix} |\downarrow\rangle_j & |\uparrow\rangle_j & 0 \\ 0 & 0 & |\downarrow\rangle_j \\ 0 & -|\downarrow\rangle_j & -\frac{f}{2}|\downarrow\rangle_j \end{pmatrix} \tag{4}$$

representing the spin state at site j. The three degenerate ground states of the local Hamiltonian $h_{j,j+1,j+2}$ with energies E_0, E_2 and E_{5-} can be expressed as the matrix $m_j m_{j+1} m_{j+2}$. In fact, the matrix elements are the linear combinations of these three eigen states. It is clear that $M_n = \Pi_{j=1}^{N} m_j$ gives the states of n spins. $h_{j,j+1,j+2}$, operates only on the product $m_j m_{j+1} m_{j+2}$ (whose elements are its eigen states,) without affecting other matrices. Hence, the operation of Hamiltonian will not alter the form of M_n and its matrix elements are linear combinations of the eigen states. We can write M_n in a more compact form

$$M_n = \begin{pmatrix} |\phi_n^a\rangle & |\phi_n^b\rangle & |\phi_n^c\rangle \\ 0 & -f_{n-1}|\phi_n^a\rangle & f_n|\phi_n^a\rangle \\ 0 & -f_n|\phi_n^a\rangle & f_{n+1}|\phi_n^a\rangle \end{pmatrix}$$

where $|\phi_n^a\rangle = |0\rangle, |\phi_n^b\rangle = \sum_{j=1}^n f_{n-j+1}|j\rangle$ and $|\phi_n^c\rangle = \sum_{j=1}^n f_{n-j}|j\rangle$, with $|0\rangle = |\downarrow\downarrow\downarrow\downarrow\ldots\rangle, |j\rangle = s_j^+|0\rangle$ and the coefficients f_j satisfying the relations $f_0 = 0, f_1 = 1, f_j + \frac{f}{2}f_{j+1} + f_{j+2} = 0$. Note that $f_j = cos(jk)$ where

$$k = cos^{-1}(\frac{-f}{4}), \tag{5}$$

satisfies above equations. The first eigen state is the fully polarized (FP) state and the other two can be expressed as the linear combination of two independent one-magnon states: $|\phi_n^{b'}\rangle = \frac{1}{\sqrt{n}}\sum_{j=1}^n e^{ikj}s_j^+|0\rangle$, $|\phi_n^{c'}\rangle = \frac{1}{\sqrt{n}}\sum_{j=1}^n e^{-ikj}s_j^+|0\rangle$ where j is the site position. The periodic boundary condition implies that $Nk = 2m\pi$. Thus we have constructed the eigen states for a zigzag spin chain of arbitrary length at the surface Eq.(2).

Proof of the quantum critical surface by positive semi-definite matrix theorem: We use the theorem of positive semi-definite matrix to show that for spin chain of arbitrary length, the above mentioned three degenerate eigen states are actually the global ground state in certain region. The theorem of positive semi-definite matrix is the following: *The necessary and sufficient condition for a real symmetric matrix A to be positive semi-definite is $x^T Ax \geq 0$ for all real vectors x. If M and N are positive semi-definite, then the sum $M+N$, and the direct sum $M \oplus N$, and direct product $M \otimes N$ are also positive semi-definite.*

The FP state and the one-magnon states are the degenerate ground states, if the matrix $(h_{j,j+1,j+2} - E_0\hat{1})$, with E_0 being their energy, is a positive semi-definite matrix. We begin the proof with a local Hamiltonian with three spins. By setting $x = \frac{f(f+4\Delta_1)}{2(2+2\Delta_2+\Delta_1 f)}$, Eq.(2) and writing the Hamiltonian $h_{i,i+1,i+2} - E_0$ in a real and symmetric matrix form, we proceed to prove it is a positive semi-definite matrix. The value $\mathbf{x}^T(h_{i,i+1,i+2} - E_0)\mathbf{x}$ where $\mathbf{x} = (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)^T$ is any real column matrix, is

$$x^T(h_{i,i+1,i+2} - E_0)x = \frac{1}{4}[4(1+\Delta_2+\Delta_1 f + \frac{f^2}{8})x_1^2 + \frac{f(4\Delta_1+f)}{2}(x_2^2+x_5^2) + (4+4\Delta_2-\frac{f^2}{2})x_3^2 + 2(x_2+\frac{f}{2}x_3+x_5)^2 + 2(x_4+\frac{f}{2}x_6+x_7)^2]. \tag{6}$$

$(h_{j,j+1,j+2} - E_0\hat{1})$ is a positive semi-definite matrix provided two inequalities below are satisfied simultaneously

$$f(4\Delta_1+f) \geq 0, \quad 4+4\Delta_2-\frac{f^2}{2} \geq 0. \tag{7}$$

Since all the local Hamiltonian $(h_{j,j+1,j+2}-E_0\hat{1})$ are positive semi-definite matrices, so is $h_{123}\otimes\hat{1}_2+\hat{1}_2\otimes h_{234}$ and $H-E_0$. Thus we conclude that in the region given by inequalities (7), the degenerate ground state is the FP state and one magnon state with k in Eq.(5). The degenerate ground states are $|\phi_N^a\rangle = |0\rangle$, $|\phi_N^b\rangle = \sum_{m=1}^N e^{ikm} s_m^+ |0\rangle$ $|\phi_N^c\rangle = \sum_{m=1}^N e^{-ikm} s_m^+ |0\rangle$. We will give argument below that even if outside of the region given by inequalities (7), they are still the degenerate ground states. In the mean time we show the quantum critical surfaces given by Eq.(2) for the case $\Delta_1=\Delta_2=d$ in Fig.1(a) and the case $\Delta_1=d$ and $\Delta_2=0$ in Fig.1(b). In the region above the surfaces only the FP state is the ground state. There are several interesting features. Firstly, larger Δ_1 and Δ_2, requires larger magnetic field b. This means that the system is made difficult to be polarized by anisotropy. On the other hand, negative f, ferromagnetic coupling, reduces the value of b as expected. Eq.(2) is valid for $f\geq-4$. Beyond which only the FP state is the ground state. Quantitatively, we see a linear dependence of b on Δ_1 and Δ_2. This indicates that treating the coupling $S_i^z S_j^z$ with the mean-field approximation is accurate at least in the region close to that given by Eq.(2).

Analytical solutions outside the region of Eq.(7)): Beyond the regions given by (7), the global Hamiltonian is still likely to be positive semi-definite. We apply Bethe ansatz to analyze this region. Our approach is the following. We write down the Bethe ansatz solution for n up-spins state, and the constraints of the coefficients. For a given b and a given f, we find the state with lowest energy. We are able to show that for $n\leq\frac{N}{3}$, the energy of above mentioned state is greater or equal to the energy of FP state on the surface given by Eq.(2). We begin by writing the state as the linear combinations of n spin-up states:

$$|\psi\rangle = \sum_{x_1<x_2<\cdots<x_n} a_{x_1,x_2,\ldots,x_n} s_{x_1}^+ s_{x_2}^+ \ldots s_{x_n}^+ |0\rangle \qquad (8)$$

where $a_{x_1,x_2,\ldots,x_n}$ is symmetric under exchanges of spin indices. Their constraints can be obtained by requiring $|\psi\rangle$ to be the eigen state of the Hamiltonian.

In this case there are many constraints. But one of them has the property $x_{i+1}>x_i+2$ with all possible i. This means that all the up-spins are separated by at least two down-spins, or no two up-spins are NN or NNN. This relation is

$$\begin{aligned}
&2(E-E_0)a_{x_1,x_2,\ldots,x_n}\\
&=2n(b-\Delta_2-\Delta_1 f)a_{x_1,x_2,\ldots,x_n}\\
&+f\sum_{i=1}^{n}(a_{\ldots x_{i-1},x_i-1,x_{i+1}\ldots}+a_{\ldots x_{i-1},,x_i+1,x_{i+1}\ldots})\\
&+\sum_{i=1}^{n}(a_{\ldots x_{i-1},x_i-2,x_{i+1}\ldots}+a_{\ldots x_{i-1},,x_i+2,x_{i+1}\ldots})
\end{aligned}$$



FIG. 1: The quantum critical surface in the case (a) $\Delta_1=\Delta_2=d$, and (b)$\Delta_1=d,\Delta_2=0$.

With the same method for two-magnon states, we let $a_{x_1,x_2,\ldots,x_n}=\sum_P A_P\exp[i\sum_{j=1}^n k_{Pj}x_j]$, where P means permutation of spin sites j. The equation becomes

$$2(E-E_0)=2n(b-\Delta_2-\Delta_1 f)+2f\sum_{i=1}^{n}\cos k_i+2\sum_{i=1}^{n}\cos 2k_i. \qquad (9)$$

If k_i's were independent variables, then the lowest energy would have been obtained by varying them independently. The result is

$$k_i=k=\pm\cos^{-1}(-\frac{f}{4}). \qquad (10)$$

For $E=E_0$ and Eq.(10), we again reach the relation in Eq. (2). Since k_i's have other constraints, the energy we thus get is the lower bound. Hence, we conclude that for $1\leq n\leq\frac{N}{3}$, the FP state and one-magnon states have the lowest energy with the magnetic field given by Eq. (2). It is plausible to assume that this is true for arbitrary value of n. Bethe ansatz gives the excitation energy of magnons as $\Delta E = E - E_0 = 1 + \frac{f^2}{8} + f\cos k + \cos 2k$. Interestingly, it does not depend on the anisotropy Δ_1 and Δ_2. It vanishes when Eq.(5) is satisfied. Physically, this means that the magnons soften at certain wave vectors, depending on the value of f. This phenomena can be detected by neutron scattering or susceptibility measurement.

Application to superconducting quantum dot system: Here we derive the analytical expression of the quantum critical surface of a superconducting quantum dots (SQD) array with NN and NNN tunneling and Coulomb interaction in the presence of gate voltage in every superconducting dot. The quantum critical surface separate the charge density wave phase of lower commensurability ($n = 1/2$), which corresponds to a anti-ferromagnetic phase of a spin chain; from a higher commensurability state ($n = 1$, or integer numbers of Cooper pair in every dot,) which correspondence to the FP phase. Therefore, we use the relation between b and f evaluated in the previous section to study the quantum critical surface between these two phases. The quantum fluctuations are controlled by the system parameters like charging energy of SQD and the Josephson coupling (E_J). In the SQD model [21], we consider the finite range E_J and Coulomb interaction which have experimental support [22–25]. The model Hamiltonian of our study consists of different interactions, $H = H_{J1} + H_{J2} + H_{EC0} + H_{EC1} + H_{EC2}$. We recast our basic Hamiltonians in the spin language around the charge degeneracy point. $H_{J1} = 2E_{J1}\sum_i({S_i}^{\dagger}{S_{i+1}}^{-} + h.c)$, $H_{J2} = 2E_{J2}\sum_i({S_i}^{\dagger}{S_{i+2}}^{-} + h.c)$, $H_{EC0} = \frac{E_{C0}}{2}\sum_i(2{S_i}^Z + h)^2 \approx 2hE_{C0}\sum_i 2{S_i}^Z$, $H_{EC1} = 4E_{Z1}\sum_i {S_i}^Z\ {S_{i+1}}^Z$, $H_{EC2} = 4E_{Z2}\sum_i {S_i}^Z\ {S_{i+2}}^Z$. where H_{J1} and H_{J2} are Josephson energy Hamiltonians respectively for NN and NNN Josephson tunneling between the SQD and H_{EC0}, H_{EC1}, and H_{EC2} are respectively the Hamiltonians for on-site, NN and NNN charging energies of the SQD. $h = \frac{2n+1-N}{2}$ is a parameter which allows tuning the system to a degeneracy point by means of gate voltage. When $E_{C0} >> E_J$, sequential tunnelling of Cooper pair across the SQD is not a energetically favorable process. One must consider the co-tunneling effect, i.e., the higher order expansion in $\frac{E_{J1}}{E_{C0}}$, which reduce the intersite Coulomb charging energy and also increase the NNN E_J [21, 26, 27]. This results in new tunneling energies: $E_{J2} \to E_{J2} + {E_{J1}}^2/2E_{C0}$, and $E_{Z1} \to E_{Z1} - 3{E_{J1}}^2/16E_{C0}$.

Now we present the analytical expression of quantum critical surface for the presence and absence of co-tunneling effect, following the relation b and f in Eq.(2). The analytical expression of quantum critical surface in the absence of co-tunneling effect are the following

$$2hE_{C0} = 2E_{J2} + 4(E_{z1} + E_{z2}) + \frac{{E_{J1}}^2}{4E_{J2}}$$

while that in the presence of co-tunneling effect is

$$2hE_{C0} = 2E_{J2} + 4(E_{z2} + E_{z1}) + \frac{E_{J1}^2}{4} + \frac{E_{J1}^2}{4E_{J2}} + \frac{2E_{J1}^2}{E_{C0}}.$$

We have already calculated the quantum critical surface on which FP state and one magnon state are degenerate. Here the one vacant site of Cooper pair in the SQD lattice array corresponds to one magnon state and Mott insulating state with integer numbers of Cooper pairs in each lattice site corresponds to the FP state. The applied gate voltage on each SQD of the array corresponds to the applied magnetic field. The charging energy corresponds to the anisotropy of the exchange interaction of the spin system. For large gate voltage, the SQD is in the insulating state. Interestingly, the charging energy tends to reverse this trend by destabilizing the state of integer numbers of Cooper pairs in each lattice site and makes the system conducting. The analytical expression for the quantum critical surface for the SQD array is the first in the literature.

Conclusions: We have found the quantum critical surface of a zigzag spin chain under magnetic field by using the theorem of positive semi-definite matrix and Bethe-ansatz methods. We have also find the quantum critical surface of superconducting quantum dots system by using our analytical methods.

[1] U. Schollwock, J. Richter, D.J.J. Farnell, R.F. Bishop (Eds.), Lect. Notes Phys. **645**, "Quantum Magnetism"(Springer, Berlin Heidelberg 2004).

[2] H. Bethe, Zeitschrift fu"r Physik A, Vol.**71**, 205 (1931).

[3] M. Karbach and G. Muller, arXiv: cond-mat/9809162 (1998),

[4] Minoru Takahashi, "Thermodynamics of one-dimensional solvable models"(Cambridge 1999).

[5] A. A. Zvyagin, J. Phys. A: Math. Gen **34**, R21 (2001)

[6] V. E. Korepin , N. M. Bogoliubov and A. G. Izergin , "Quantum Inverse Scattering Method and Correlation Functions"(Cambridge 1993).

[7] A, Kl*ü*mper, A. Schadschneider, and J. Zittartz, J. Phys. A **24**, L955 (1995); A, Kl*ü*mper, A. Schadschneider, and J. Zittartz, Z. Phys: B -Cond. Mat. **87**, 281 (1992).

[8] K. Totsuka and M. Suzuki, J. Phys. C 1639 (1995),

[9] A. K. Kolezhuk, Phys. Rev. B **49**, 8901 (1994),

[10] Gang Su, Phys. Lett. A **213**, 93 (1996),

[11] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).

[12] C. K. Majumdar, D. k. Ghosh, J. Math. Phys. **10**,1388 (1969).

[13] T. Hanada, J. Kane, S. Nakagawa and Y. Natsume, J. Phys. Soc. Japan **57**, 1891 (1988).

[14] Steve R. White, Phys. Rev. B **54**, 9862 (2008),

[15] D. Allen and D. S*é*n*é*chal, Phys. Rev. B **51**, 6394 (1995).

[16] S. Rao and D. Sen, J. Phys.: Condens. Matter **9**, 1831 (1997).

[17] S. Sarkar and D. Sen, Phys. Rev. B **65**, 172408 (2002),

[18] Y. Mizuno, T. Tohyama and S. Mackawa, Phys. Rev. B **57**, 5326 (1998).

[19] H. Eisaki, N. Motoyama, and S. Uchida, Physica C **282**, 1323 (1997).

[20] D. C. Cabra, A. Honecker and P. Pujol, Eur. Phys. J. B **13**, 55 (2000).

[21] S. Sarkar, Phys. Rev. B **75**, 014528 (2007).

[22] H. M. Jaeger *et al.*, Phys. Rev. B **40**, 182 (1989).

[23] H. S. J. van der Zant *et al.*, Phys. Rev. Lett, **69**, 2971 (1992).

[24] C. D. Chen *et al.*, Phys. Rev. B **51**, 15645 (1995).

[25] E. Chow, Phys. Rev. Lett. **81**, 204 (1998).

[26] L. I. Glazman and A. I. Larkin, Phys. Rev. Lett **79**, 3736 (1997).

[27] D. Giuliano and P. Sodano, Nucl. Phys. B **711**, 480 (2005).